Neuroscience Letters, 156 (1993) 176–178
Elsevier Scientific Publishers Ireland Ltd.

AUTHOR INDEX

Aguilar-Roblero, R., see Zainos-Rosales, A., 99
Aigouy, L., see Cadet, R., 43
Aizawa, T., see Hisano, S., 21
Akiguchi, I., see Nagao, M., 47
Akiguchi, I., see Tomimoto, H., 105
Albala, N., see Mailleux, P., 125
Albe-Fessard, D., see Marcilloux, J.-C., 113
Alcázar, A., see Martín, M.E., 109

Baudrie, V., Oliver, C. and Chaouloff, F., Corticosterone response to the serotonergic agonist D-fenfluramine may be independent from corticotropin-releasing factor (CRF), 121
Bekkers, J.M. and Stevens, C.F., NMDA receptors at excitatory synapses in the hippocampus: test of a theory of magnesium block, 73
Ben-Ari, Y., see Represa, A., 149
Bernet, F., see Maubert, E., 5
Boer, G.J., Snijdewint, F.G.M., Licht, R. and te Riele, P., Increased loss of brain DNA in the neonatal vasopressin-deficient Brattleboro rat, but not in normal rat treated with vasopressin antagonist, 17
Brinkmeier, H., Wollinsky, K.H., Seewald, M.J., Hülser, P.-J., Mehrkens, H.-H., Kornhuber, H.H. and Rüdel, R., Factors in the cerebrospinal fluid of multiple sclerosis patients interfering with voltage-dependent sodium channels, 172
Bugnon, C., see Griffond, B., 91

Cabanis, E.A., see Marcilloux, J.-C., 113
Cadet, R., Aigouy, L. and Woda, A., Sustained hyperalgesia can be induced in the rat by a single formalin injection and depends on the initial nociceptive inputs, 43
Cammarata, S., Caponnetto, C. and Tabaton, M., Ubiquitin-reactive neurites in cerebral cortex of subjects with Huntington's chorea: a pathological correlate of dementia?, 96
Caponnetto, C., see Cammarata, S., 96
Castoldi, A.F., see Tan, X., 13
Chaouloff, F., see Baudrie, V., 121
Chikamori-Aoyama, M., see Hisano, S., 21
Cimino, M., see Stankov, B., 131
Ciofi, P., see Griffond, B., 91
Clementi, F., see Stankov, B., 131
Costa, L.G., see Tan, X., 13
Croix, D., see Griffond, B., 91

Deray, A., see Griffond, B., 91
Dooley, D.J., see Ginap, T., 35
Drucker-Colín, R., see Zainos-Rosales, A., 99
Dunbar, J.S., see Inglis, W.L., 78
Dupouy, J.P., see Maubert, E., 5

Fando, J.L., see Martín, M.E., 109
Félix, M.-B., see Marcilloux, J.-C., 113
Fellmann, D., see Griffond, B., 91
Feuerstein, T.J., see Ginap, T., 35
Fischer-Colbrie, R., see Mahata, S.K., 70
Fraschini, F., see Stankov, B., 131
Fukui, Y., see Hisano, S., 21
Furiya, Y., Sahara, N. and Mori, H., Okadaic acid enhances abnormal phosphorylation on tau proteins, 67

García, A.M., see Martín, M.E., 109
Ghilini, G., see Represa, A., 149
Ginap, T., Dooley, D.J. and Feuerstein, T.J., The non-dihydropyridine L-type voltage-sensitive calcium channel activator FPL 64176 enhances K^+-evoked efflux of [^{3}H]norepinephrine from rat neocortical slices, 35
Girdlestone, D., see Maldonado, R., 135
Griffond, B., Deray, A., Fellmann, D., Ciofi, P., Croix, D. and Bugnon, C., Colocalization of prolactin- and dynorphin-like substances in a neuronal population of the rat lateral hypothalamus, 91

Henriksen, S.J., see Prospéro-García, O., 158
Hicks, K.J., see Hisanaga, K., 57
Hiroi, N. and White, N.M., The ventral pallidum area is involved in the acquisition but not expression of the amphetamine conditioned place preference, 9
Hisanaga, K., Sagar, S.M., Koistinaho, J., Hicks, K.J. and Sharp, F.R., VIP-induced stellation and immediate early gene expression in astrocytes: effects of dexamethasone, 57
Hisano, S., Chikamori-Aoyama, M., Aizawa, T. and Fukui, Y., Somatostatin-like immunoreactive axon terminals on oxytocin-like immunoreactive neurons in the paraventricular nucleus of the rat hypothalamus, 21
Höllt, V., see Ziólkowska, B., 39
Hömberg, V., see Ziemann, U., 167
Hülser, P.-J., see Brinkmeier, H., 172

Ibazizen, M.-T., see Marcilloux, J.-C., 113
Ikarashi, Y., see Takahashi, A., 54
Inglis, W.L., Dunbar, J.S. and Winn, P., Barbiturate anaesthesia reduces the neurotoxic effects of quinolinate but not ibotenate in the rat pedunculopontine tegmental nucleus, 78
Ishimaru, H., see Takahashi, A., 54

Janáky, R., Varga, V., Saransaari, P. and Oja, S.S., Glutathione modulates the *N*-methyl-D-aspartate receptor-activated calcium influx into cultured rat cerebellar granule cells, 153
Jouvet, M., see Roky, R., 117

Kamo, H., see Nagao, M., 47
Khrestchatisky, M., see Represa, A., 149
Kimura, J., see Tomimoto, H., 105
Kimura, J., see Nagao, M., 47
Koistinaho, J., see Hisanaga, K., 57
Kornhuber, H.H., see Brinkmeier, H., 172

Laplace, J.-P., see Marcilloux, J.-C., 113
Licht, R., see Boer, G.J., 17
Lönnberg, P., see Päivärinta, M.A., 1
Lu, Q., see Wood, J.G., 83
Lucini, V., see Stankov, B., 131

Mahata, M., see Mahata, S.K., 70
Mahata, S.K., Mahata, M., Fischer-Colbrie, R. and Winkler, H., Vesicle monoamine transporters 1 and 2: differential distribution and regulation of their mRNAs in chromaffin and ganglion cells of rat adrenal medulla, 70

Neuroscience Letters, 156 (1993) 176–178
Elsevier Scientific Publishers Ireland Ltd.

AUTHOR INDEX

Aguilar-Roblero, R., see Zainos-Rosales, A., 99
Aigouy, L., see Cadet, R., 43
Aizawa, T., see Hisano, S., 21
Akiguchi, I., see Nagao, M., 47
Akiguchi, I., see Tomimoto, H., 105
Albala, N., see Mailleux, P., 125
Albe-Fessard, D., see Marcilloux, J.-C., 113
Alcázar, A., see Martín, M.E., 109

Baudrie, V., Oliver, C. and Chaouloff, F., Corticosterone response to the serotonergic agonist D-fenfluramine may be independent from corticotropin-releasing factor (CRF), 121
Bekkers, J.M. and Stevens, C.F., NMDA receptors at excitatory synapses in the hippocampus: test of a theory of magnesium block, 73
Ben-Ari, Y., see Represa, A., 149
Bernet, F., see Maubert, E., 5
Boer, G.J., Snijdewint, F.G.M., Licht, R. and te Riele, P., Increased loss of brain DNA in the neonatal vasopressin-deficient Brattleboro rat, but not in normal rat treated with vasopressin antagonist, 17
Brinkmeier, H., Wollinsky, K.H., Seewald, M.J., Hülser, P.-J., Mehrkens, H.-H., Kornhuber, H.H. and Rüdel, R., Factors in the cerebrospinal fluid of multiple sclerosis patients interfering with voltage-dependent sodium channels, 172
Bugnon, C., see Griffond, B., 91

Cabanis, E.A., see Marcilloux, J.-C., 113
Cadet, R., Aigouy, L. and Woda, A., Sustained hyperalgesia can be induced in the rat by a single formalin injection and depends on the initial nociceptive inputs, 43
Cammarata, S., Caponnetto, C. and Tabaton, M., Ubiquitin-reactive neurites in cerebral cortex of subjects with Huntington's chorea: a pathological correlate of dementia?, 96
Caponnetto, C., see Cammarata, S., 96
Castoldi, A.F., see Tan, X., 13
Chaouloff, F., see Baudrie, V., 121
Chikamori-Aoyama, M., see Hisano, S., 21
Cimino, M., see Stankov, B., 131
Ciofi, P., see Griffond, B., 91
Clementi, F., see Stankov, B., 131
Costa, L.G., see Tan, X., 13
Croix, D., see Griffond, B., 91

Deray, A., see Griffond, B., 91
Dooley, D.J., see Ginap, T., 35
Drucker-Colín, R., see Zainos-Rosales, A., 99
Dunbar, J.S., see Inglis, W.L., 78
Dupouy, J.P., see Maubert, E., 5

Fando, J.L., see Martín, M.E., 109
Félix, M.-B., see Marcilloux, J.-C., 113
Fellmann, D., see Griffond, B., 91
Feuerstein, T.J., see Ginap, T., 35
Fischer-Colbrie, R., see Mahata, S.K., 70
Fraschini, F., see Stankov, B., 131
Fukui, Y., see Hisano, S., 21
Furiya, Y., Sahara, N. and Mori, H., Okadaic acid enhances abnormal phosphorylation on tau proteins, 67

García, A.M., see Martín, M.E., 109
Ghilini, G., see Represa, A., 149
Ginap, T., Dooley, D.J. and Feuerstein, T.J., The non-dihydropyridine L-type voltage-sensitive calcium channel activator FPL 64176 enhances K^+-evoked efflux of [^{3}H]norepinephrine from rat neocortical slices, 35
Girdlestone, D., see Maldonado, R., 135
Griffond, B., Deray, A., Fellmann, D., Ciofi, P., Croix, D. and Bugnon, C., Colocalization of prolactin- and dynorphin-like substances in a neuronal population of the rat lateral hypothalamus, 91

Henriksen, S.J., see Prospéro-García, O., 158
Hicks, K.J., see Hisanaga, K., 57
Hiroi, N. and White, N.M., The ventral pallidum area is involved in the acquisition but not expression of the amphetamine conditioned place preference, 9
Hisanaga, K., Sagar, S.M., Koistinaho, J., Hicks, K.J. and Sharp, F.R., VIP-induced stellation and immediate early gene expression in astrocytes: effects of dexamethasone, 57
Hisano, S., Chikamori-Aoyama, M., Aizawa, T. and Fukui, Y., Somatostatin-like immunoreactive axon terminals on oxytocin-like immunoreactive neurons in the paraventricular nucleus of the rat hypothalamus, 21
Höllt, V., see Ziólkowska, B., 39
Hömberg, V., see Ziemann, U., 167
Hülser, P.-J., see Brinkmeier, H., 172

Ibazizen, M.-T., see Marcilloux, J.-C., 113
Ikarashi, Y., see Takahashi, A., 54
Inglis, W.L., Dunbar, J.S. and Winn, P., Barbiturate anaesthesia reduces the neurotoxic effects of quinolinate but not ibotenate in the rat pedunculopontine tegmental nucleus, 78
Ishimaru, H., see Takahashi, A., 54

Janáky, R., Varga, V., Saransaari, P. and Oja, S.S., Glutathione modulates the *N*-methyl-D-aspartate receptor-activated calcium influx into cultured rat cerebellar granule cells, 153
Jouvet, M., see Roky, R., 117

Kamo, H., see Nagao, M., 47
Khrestchatisky, M., see Represa, A., 149
Kimura, J., see Tomimoto, H., 105
Kimura, J., see Nagao, M., 47
Koistinaho, J., see Hisanaga, K., 57
Kornhuber, H.H., see Brinkmeier, H., 172

Laplace, J.-P., see Marcilloux, J.-C., 113
Licht, R., see Boer, G.J., 17
Lönnberg, P., see Päivärinta, M.A., 1
Lu, Q., see Wood, J.G., 83
Lucini, V., see Stankov, B., 131

Mahata, M., see Mahata, S.K., 70
Mahata, S.K., Mahata, M., Fischer-Colbrie, R. and Winkler, H., Vesicle monoamine transporters 1 and 2: differential distribution and regulation of their mRNAs in chromaffin and ganglion cells of rat adrenal medulla, 70

Mailleux, P. and Vanderhaeghen, J.-J., Glucocorticoid regulation of cannabinoid receptor messenger RNA levels in the rat caudate-putamen. An in situ hybridization study, 51
Mailleux, P., Albala, N. and Vanderhaeghen, J.-J., Ontogeny of gene expression of inositol 1,4,5-trisphosphate receptor in the rat brain: high mRNA levels in the cerebellar Purkinje cells, 125
Maldonado, R., Girdlestone, D. and Roques, B.P., RP 67580, a selective antagonist of neurokinin-1 receptors, modifies some of the naloxone-precipitated morphine withdrawal signs in rats, 135
Manzo, L., see Tan, X., 13
Marcilloux, J.-C., Félix, M.-B., Rampin, O., Stoffels, C., Ibazizen, M.T., Cabanis, E.A., Laplace, J.-P. and Albe-Fessard, D., Preliminary results of a magnetic resonance imaging (MRI) study of the pig brain placed in stereotaxic conditions, 113
Marini, P., see Stankov, B., 131
Martín, M.E., Alcázar, A., Fando, J.L., García, A.M. and Salinas, M., Translational initiation factor eIF-2 subcellular levels and phosphorylation status in the developing rat brain, 109
Marttila, R.J., see Päivärinta, M.A., 1
Maruyama, Y., see Takahashi, A., 54
Mathes, C., see Wang, S.S.-H., 145
Maubert, E., Dupouy, J.P. and Bernet, F., Effect of adrenal demedullation on neuropeptide Y content of the capsule/glomerulosa zone of the rat adrenal gland, 5
McCarron, R.M., Wang, L., Stanimirovic, D.B. and Spatz, M., Endothelin induction of adhesion molecule expression on human brain microvascular endothelial cells, 31
Mehrkens, H.-H., see Brinkmeier, H., 172
Mendoza-Ramírez, J.-L., see Zainos-Rosales, A., 99
Miller, D.R., see Prospéro-García, O., 158
Morán, J., see Sánchez-Olea, R., 141
Moreau, J., see Represa, A., 149
Mori, H., see Furiya, Y., 67
Morrisett, R.A., see Zhang, G., 27
Motoike, T., Sano, K., Nakamura, H. and Takai, Y., Expression of *smg* p25A/*rab* 3A guanine nucleotide dissociation inhibitor (GDI) in neurons and glial cells from rat brain, 87

Nagao, M., Kamo, H., Akiguchi, I. and Kimura, J., Cycloheximide downsynthesizes inhibitory molecules for neurite outgrowth in neural transplantation, 47
Nakamura, H., see Motoike, T., 87
Nakamura, S., see Tomimoto, H., 105
Netz, J., see Ziemann, U., 167

Ogawa, K., see Taniguchi, K., 24
Oja, S.S., see Janáky, R., 153
Okamura, M., see Taniguchi, K., 24
Oliver, C., see Baudrie, V., 121

Päivärinta, M.A., Marttila, R.J., Lönnberg, P. and Rinne, U.K., Decreased raphe serotonin in rabbits with experimental herpes simplex encephalitis, 1
Pasantes-Morales, H., see Sánchez-Olea, R., 141
Peña, C., see Sánchez-Olea, R., 141
Pollard, H., see Represa, A., 149
Priestley, J.V., see Rattray, M., 163
Prospéro-García, O., Miller, D.R. and Henriksen, S.J., Hippocampal interneuron activity in unanesthetized rats: relationship to the sleep-wake cycle, 158

Rampin, O., see Marcilloux, J.-C., 113
Rattray, M. and Priestley, J.V., Differential expression of GABA transporter-1 messenger RNA in subpopulations of GABA neurones, 163
Reich, C., see Wood, J.G., 83
Represa, A., Pollard, H., Moreau, J., Ghilini, G., Khrestchatisky, M. and Ben-Ari, Y., Mossy fiber sprouting in epileptic rats is associated with a transient increased expression of α-tubulin, 149
Rinne, U.K., see Päivärinta, M.A., 1
Roky, R., Valatx, J.-L. and Jouvet, M., Effect of prolactin on the sleep–wake cycle in the rat, 117
Roques, B.P., see Maldonado, R., 135
Rüdel, R., see Brinkmeier, H., 172

Sagar, S.M., see Hisanaga, K., 57
Sahara, N., see Furiya, Y., 67
Saito, H., see Taniguchi, K., 24
Salinas, M., see Martín, M.E., 109
Sánchez-Olea, R., Peña, C., Morán, J. and Pasantes-Morales, H., Inhibition of volume regulation and efflux of osmoregulatory amino acids by blockers of Cl^- transport in cultured astrocytes, 141
Sano, K., see Motoike, T., 87
Saransaari, P., see Janáky, R., 153
Scharfman, H.E., Activation of dentate hilar neurons by stimulation of the fimbria in rat hippocampal slices, 61
Seewald, M.J., see Brinkmeier, H., 172
Sharp, F.R., see Hisanaga, K., 57
Snijdewint, F.G.M., see Boer, G.J., 17
Spatz, M., see McCarron, R.M., 31
Stanimirovic, D.B., see McCarron, R.M., 31
Stankov, B., Cimino, M., Marini, P., Lucini, V., Fraschini, F. and Clementi, F., Identification and functional significance of nicotinic cholinergic receptors in the rat pineal gland, 131
Stevens, C.F., see Bekkers, J.M., 73
Stoffels, C., see Marcilloux, J.-C., 113
Szelényi, A., see Ziemann, U., 167

Tabaton, M., see Cammarata, S., 96
Takahashi, A., Ishimaru, H., Ikarashi, Y. and Maruyama, Y., Intraventricular injection of neostigmine increases dopaminergic and noradrenergic nerve activities: hyperglycemic effects and neurotransmitters in the hypothalamus, 54
Takai, Y., see Motoike, T., 87
Tan, X., Castoldi, A.F., Manzo, L. and Costa, L.G., Interaction of ethanol with muscarinic receptor-stimulated phosphoinositide metabolism during the brain growth spurt in the rat: role of acetaldehyde, 13
Taniguchi, K., Saito, H., Okamura, M. and Ogawa, K., Immunohistochemical demonstration of protein gene product 9.5 (PGP 9.5) in the primary olfactory system of the rat, 24
te Riele, P., see Boer, G.J., 17
Thompson, S.H., see Wang, S.S.-H., 145
Tomimoto, H., Akiguchi, I., Wakita, H., Nakamura, S. and Kimura, J., Distribution of NADPH-diaphorase in the cerebral blood vessels of rats: a histochemical study, 105

Valatx, J.-L., see Roky, R., 117
Vanderhaeghen, J.-J., see Mailleux, P., 51, 125
Varga, V., see Janáky, R., 153

Wakita, H., see Tomimoto, H., 105
Wang, L., see McCarron, R.M., 31
Wang, S.S.-H., Mathes, C. and Thompson, S.H., Membrane toxicity of

the protein kinase C inhibitor calphostin A by a free-radical mechanism, 145
White, N.M., see Hiroi, N., 9
Winkler, H., see Mahata, S.K., 70
Winn, P., see Inglis, W.L., 78
Woda, A., see Cadet, R., 43
Wollinsky, K.H., see Brinkmeier, H., 172
Wood, J.G., Lu, Q., Reich, C. and Zinsmeister, P., Proline-directed kinase systems in Alzheimer's disease pathology, 83

Zainos-Rosales, A., Aguilar-Roblero, R., Mendoza-Ramírez, J.-L. and Drucker-Colín, R., 2-Deoxyglucose autoradiographic evidence that fetal substantia nigra grafts exert contralateral effects in 6-hydroxydopamine-lesioned animals who recover motor asymmetries, 99
Zhang, G. and Morrisett, R.A., Ethanol inhibits tetraethylammonium chloride-induced synaptic plasticity in area CA1 of rat hippocampus, 27
Ziemann, U., Netz, J., Szelényi, A. and Hömberg, V., Spinal and supraspinal mechanisms contribute to the silent period in the contracting soleus muscle after transcranial magnetic stimulation of human motor cortex, 167
Zinsmeister, P., see Wood, J.G., 83
Ziólkowska, B. and Höllt, V., The NMDA receptor antagonist MK-801 markedly reduces the induction of c-fos gene by haloperidol in the mouse striatum, 39